


HEIGHT ADJUSTABLE PLATFORMS



Picture Sources – Varidesk, Ergotron



CSU Ergonomics Program
Office of Risk Management & Insurance
141 General Services Building
Fort Collins, CO 80523-6002
rmi.prep.colostate.edu/ergonomics



Height Adjustable – Sit/Stand Workstations

We have all heard it; sitting is the new smoking. Being sedentary shortens our lifespan and can create long-term negative health effects. These negative health outcomes include all-cause mortality (van der Ploeg, Chey, Korda, Banks, & Bauman, 2012) disabilities such as obesity, diabetes, and cardiovascular disease (MacEwen, MacDonald, & Burr, 2015; Wilmot et al., 2012) as well as musculoskeletal disorders (MSD's) of the hand, wrist, neck, upper and lower back (Ekman, Andersson, Hagberg, & Heljm, 2000; Gerr et al.; 2002; Korhenon et al.; 2003; Rocha et al., 2005; Wahlström, 2005; Wahlström, Hagberg, Toomingas, & Tornqvist, 2004).

Although sitting for extended periods can certainly be detrimental, standing for extended periods is not the answer. Dr. Jack Dennerlein, Adjunct Professor of Ergonomics and Safety in the School of Public Health at Harvard University has said, *“Everybody says sitting is the new smoking. You have to remember, standing is the old smoking”* (Ducharme, 2016). Detriments of prolonged standing can include malfunction of venous valves, venous insufficiency, etc. (Krijnen, de Boer, Ader, & Bruynzeel, 1997), have a detrimental effect on arterial blood flow (Reinhardt et al., 2000) and increase the risk of carotid atherosclerosis (Krause et al. (2000). Extended durations of standing can also lead to long-lasting muscle fatigue, which may be present without being noticed and this may contribute to musculoskeletal disorders and back pain (HFES, 2015). Standing too much can also lead to pain in the legs, knees and feet (Neuhaus et al., 2014) and in laboratory studies, standing was perceived as less comfortable and more fatiguing than sitting (Beers, Roemmich, Epstein, & Horvath, 2008).

Neither sitting nor standing for long periods are good for the body. The key with any height adjustable workstation is movement. Unfortunately, too much focus has been placed on standing more and sitting less, when the mixture of the two postures is most important. Ideally, when using a sit/stand height adjustable workstation, users should adjust and vary postures every 30-60 minutes and avoid long durations of either sitting *or* standing (Callaghan, 2015). In contrast, Dr. Alan Hedge (2015) has recommended 20 minutes of sitting, 8 minutes of standing, and 2 minutes of stretching. Although more studies are needed to determine exact durations we should sit or stand, remember, the best posture is the next posture.

Even if you have an adjustable workstation, it does not mean you are free from all ergonomic risk. Dr. Hedge (2016) has also stated, *“So called ergonomic products do not guarantee healthy work posture!”* The correct height of the table, height and position of the keyboard and mouse, monitor(s), etc., are crucial regardless of whether sitting or standing. In a study by Lin, Catalano and Dennerlein (2016), it was found that study participants adjusted the placement of desktop height, monitor height/tilt and the keyboard distance when comparing sitting to standing. Ultimately, this means that the monitor,

Although height adjustable, 'platforms' which sit on the desktop are too high for **~95% of users**. Forcing the work too high increases likelihood for awkward postures which increases the risk for injury.

keyboard/mouse and desktop height need to be properly adjusted when alternating between sitting and standing. Simply raising or lowering the desktop height is not enough. Other components of the workstation need appropriate adjustment as well. Unfortunately, these adjustments are often neglected.

Although the ability to stand and work on the computer can be beneficial, making modifications to the workstation to allow for this should not introduce ergonomic hazards into the workstation. This can unfortunately be a common problem given the highly


available and relatively inexpensive ‘platforms’ which are attached to or sit on top of an existing table or work surface. Although desktop ‘platforms’ may be height adjustable, there are significant detriments introduced into the workstation which can expose the user to ergonomic hazards and increase the risk for injury. One of the main issues with ‘platforms’ that are placed on a fixed work surface are that these platforms do not adjust appropriately below a standard work surface height. This is crucial for a majority of the population.



Picture Source - Varidesk

The standard work surface height, which was designed for the 95th-99th percentile male (~6’2-6’4”) is approximately 29-31 inches high. This includes furniture from major furniture manufacturers or built in desk systems you might see in a reception area that have an appealing aesthetic design like a granite surface. Much like the design of a doorway, which is designed to allow people to fit underneath or inside the frame without hitting one’s head, this work surface height allows for a majority of the population to fit underneath the surface without encountering leg obstructions (knees/thighs bumping the work surface). Although this height allows most users to fit underneath the surface, it does not allow for an ideal working elbow height for approximately 95% of workers.

To further illustrate, think of the work surface height related to shoe size. Much as a shoe should fit the foot, the work surface, desk or table should adjust to ‘fit’ the body. For an ideal ‘fit’, the height of the work surface (i.e. height of the keyboard and mouse) should be located at or slightly below resting elbow height (see picture at right). When the work is designed to be performed at elbow height, the risk for shoulder, neck and back strain and fatigue due to awkward postures is minimized.



Regardless of the type of height adjustable work surface option used, if the surface does not adjust properly, (below the standard 29-31 inch work surface height), it will force the work too high. This will be the case for anyone shorter than approximately 6’2” tall (or approximately 95% of the working population).

To further illustrate using the shoe size example, using a standard ‘fixed’ work surface height of ~29 inches is like wearing a 12-14 size shoe and expecting it to ‘fit’ properly. Of course, not many individuals correctly fit into this size of shoe which is far too big for the foot. Now, think of wearing this size of shoe while running marathon. Although a person may be able to complete the marathon, it is likely that not only will their performance suffer but so will their feet, especially over the long term.

If a height adjustable ‘platform’ is added to a standard 29-31 inch ‘fixed height’ work surface (and assuming the platform does not allow for adjustment below the work surface), the keyboard and mouse are raised even higher (~1 inch). When sitting, this immediately exposes the user to awkward postures and other ergonomic risk factors which can over time lead to injury. Back to the shoe size example, now with this platform added to the work surface, the user is wearing an even larger shoe size of ~13-15 plus.

Ultimately, platforms that sit on top of a standard fixed work surface height simply will not adjust to ‘fit’ most individuals correctly. If used, exposure to ergonomic injury risk factors is increased, as is the risk for injury. Height adjustable options need to lower appropriately (beneath 29 inches) for ~95% of users,


not higher. An ideal height adjustment range which will allow for a majority of workers to adjust the work surface height to properly 'fit' whether sitting or standing is ~22-48 inches (Eastman Kodak, 2004; HFES, 2007; Sanders, 1993; BIFMA 2013; Pheasant, 1986; Konz & Johnson, 2007).

Adjusting from 22-50" would be even better and allow for taller users to stand at an appropriate height. If this specification is met, a majority of the work force will be able to adjust the work surface height to allow for the proper fit.

Desktop Height Illustration

The below illustration shows the difference in work surface height for a 95th percentile male (~6'2") compared to a 5th percentile female (~5'1"). This ~7-inch difference is significant and commonly leads to awkward postures. These detrimental awkward postures are further discussed below.

Even for an average 5'8" male (not shown) the work surface is ~4 inches too high. (An ideal work surface height for 5'8" male is approximately 26-27 inches). We might think that the difference is negligible but again, much like wearing a shoe that is 4 sizes too big, over the long term, it is likely the body will be negatively affected.



The Computer Hunch

Because the standard 29-31 inch work surface forces the keyboard and mouse too high, very common awkward postures are assumed in the neck, shoulders, elbows and upper back. As illustrated in the picture at right, leaning forward and 'hunching' over the work surface increases the lumbar disc pressure by 85% in comparison to standing in a relaxed posture (Wilke, H., Neef, P., Caimi, M., et al., 1993). Along with slouching or hunching over the keyboard and mouse, the shoulders are also affected and often round forward, elevate and or abduct (swing out to the side). These awkward postures increase tension and fatigue on the shoulders and upper back which can lead to fatigue and discomfort and over time, can lead to injury.



Picture Source: Humanscale Ergonomics

Sitting at a computer and hunching over the desktop increases back pressure by 85% compared to standing upright.

If any components of the workstation are neglected, the risk for musculoskeletal disorders can increase. Studies have shown that musculoskeletal disorders seen in computer users (i.e. tendonitis, carpal tunnel, neck strain, etc.) can be correlated to common ergonomic injury risk factors such as awkward posture, force, repetition, and duration. (Wahlstrom, 2005).

Standing Height

Work surface height is not only important when sitting. If the work surface does not adjust sufficiently when standing, similar ergonomic hazards can be introduced and can be equally damaging to the body. The increase in ergonomic risk and exposure to ergonomic hazards are commonly seen with taller individuals because the work surface does not rise adequately. Height adjustable 'platforms' may not adjust high enough to 'fit' taller users due to a limited height adjustment range. This will increase ergonomic injury risk due to a work surface that is too low. These ergonomic risk factors include extension of the elbow and wrist, rounding of the shoulders, and back flexion (leaning forward at the waist). Back to the shoe example, the taller user is being asked to wear a size 8-9 shoe when a size 12-14 is needed. Again, improper fit.

Due to these and other several other factors, 'platforms' which sit on an existing desktop are not recommended as they are not ergonomic options, do not provide a proper fit for ~90-95% of the working population and expose the user to ergonomic hazards which increase the risk for injury.

As already mentioned, an electric height adjustable table that adjusts from ~22-48 inches ensures that the work surface has the capability to adjust to properly fit a majority of the population. A table which meets ergonomic design criteria will properly 'fit' employees regardless of their height, weight, gender, etc. An electric height adjustable table, which meets these design criteria, is the truly 'ergonomic' option and has no significant limitations.



Picture Source: Herman Miller

Responsibility then belongs to the user to ensure proper adjustments are made to the work surface height and other components of the workstation in order to reduce the exposure to ergonomic hazards, minimize injury risks and decrease the likelihood for injury. However, it is not enough to simply provide adjustable workstations and expect users to adjust them correctly. Training is needed. The effectiveness of height adjustable workstations is improved with proper ergonomics training (Robertson, Ciriello, & Barabet, 2013). In addition, new technology, which is built into the height adjustable workstation itself or as part of a software program, may help with educating users regarding the proper height adjustment and appropriate workstation setup.

OSHA, Viewing Height and Distance

As mentioned above, the monitor height and tilt need to adjust relative to desk height whether sitting or standing. Proper monitor adjustment will help prevent issues such as eye strain, computer vision syndrome, awkward neck postures and head and neck discomfort. OSHA recommends a monitor viewing distance of ~20-40 from the eyes when on the computer. Many desktop 'platforms' will not allow for this distance because of the lack of space on the surface of the platform or simply because the monitors cannot at all be adjusted in depth because they are affixed to an 'arm or pole'. For this reason, these options do not meet OSHA guidelines and are not recommended.

Along with viewing distance, viewing height may also be an issue. Because certain 'platforms' force monitors to attach to an 'arm' or 'pole', many have a limited height range adjustment. If a user needs a slightly higher or lower monitor height, this adjustment simply cannot be made with many of the platform options on the market. Monitor height and depth is especially important if a user has corrective lenses (i.e. multifocal lenses). Most multifocal users need the monitor lower than eye height and many of these 'platforms' simply do not allow for the appropriate adjustments.

Cost

Although historically the cost of a height adjustable table was thought to be very expensive, height adjustable tables today may cost **less than** the average priced ergonomic chair. For additional information and a comparison between various height adjustable options, go to the below website.

[Comparison Between Height Adjustable Workstations](#)

Certainly, the cost of a height adjustable 'platform' can be relatively inexpensive in comparison to other options; however, what is the user sacrificing by adding this less than ideal option? Will it adjust to allow for a proper 'fit'? Will this expose the user to awkward postures (hands, wrists, elbows, shoulders, neck)? Is the user willing to risk an injury? Is this lack of adjustment worth the money savings? Is standing at all worth the risk?

According to Dunstan et al. (2012) 1-2 minutes of movement per hour mitigates the detrimental health effects of prolonged sitting. Additionally, at the National Ergonomics Conference in November 2015, Dr.

Joan Vernikos (former Director of Life Sciences at NASA), simply said to stand up often. “Standing up often, at least 30 times a day, is a powerful antidote to prolonged sitting.” Dr. Vernikos also mentioned that it is the frequency or number of times one stands, not the duration, that is important. Furthermore, if you cannot stand, move more often (instead of sitting for long periods and taking long breaks) as this has been shown to have a beneficial impact (Owen, Healy, Matthews, & Dunstan, 2010).

Rather than increasing the risk for injury with the use of a ‘platform’ that does not properly fit the body, stand up frequently. Get a refill of water or coffee, walk to the printer, get up and walk to a co-worker’s office instead of sending an email, get up to stretch, get a telephone headset and stand while taking phone calls, arrange walking meetings, park further away from your building, take the stairs, or leave your office for lunch. Remember, movement throughout the day is crucial.

In summary, ergonomics is about improving human performance. With the proper fit of equipment to each individual, human performance will improve. Rather than invest in ‘platforms’ which have limitations outlined in this document, invest in a properly designed height adjustable table and work safely and comfortably.

Additional Information Height Adjustable Workstations

- [Cornell University – Sit Stand](#)
- [Cornell University – Sit Stand Programs](#)
- [Workrite Ergonomics The Facts About Standing Desks White Paper](#)

Ergonomics Program Resources

- [CSU Ergonomics Program Home Webpage](#)
- [Ergonomic Evaluation Request](#)
- [Ergo Lab & Equipment Trials](#)
- [Ergonomics Matching Funds Program](#)

Contact Information

FRANK GONZALES, CPE
ERGONOMICS ADMINISTRATOR

Tel 970-491-2724

Fax 970-491-4804

Frank.Gonzales@colostate.edu

Subject to change without notice.

References

- Beers, E., Roemmich, J., Epstein, L., Horvath, P. (2008) Increasing passive energy expenditure during clerical work. *Eur J Appl Physiol* (2008) 103:353–360
- Davis, K., & Kotowski, S. (2014). Postural Variability. *Human Factors: The Journal of Human Factors and Ergonomics Society*, 56(7), 1249-1261.
- Ducharme, J. (2016). Boston Magazine. Six Tips for Using Standing Desks Correctly. Retrieved on November 10, 2016, from <http://www.bostonmagazine.com/health/blog/2016/05/10/standing-desks/>
- Dunstan, D., Kingwell, B., Larsen, R., Healy, G., Cerin, E., Hamilton, M., Shaw, J., Bertovic, David A., Zimmet, P., Salmon, J., Owen, N. (2012). Breaking Up Prolonged Sitting Reduces Postprandial Glucose and Insulin Responses. *Diabetes Care*, 35(5), 976-983.
- Gerr, F., Marcus, M., Monteilh, C., Ortiz, D., Gentry, E., Cohen, S., Edwards, A., Ensor, C., Kleinbaum, D., (2002). A prospective study of computer users: II. Postural risk factors for musculoskeletal symptoms and disorders. *American Journal of Industrial Medicine*, 41(4), 236-249.
- Grimby-Ekman, A., Andersson, E., & Hagberg, M. (2009). Analyzing musculoskeletal neck pain, measured as present pain and periods of pain, with three different regression models: A cohort study. *Bmc Musculoskeletal Disorders*, 10, *Bmc Musculoskeletal Disorders*, 2009 Jun 19, Vol.10.
- Eastman Kodak Company, *Ergonomic Design for People at Work*, 2nd ed., John Wiley & Sons, Inc., Hoboken, NJ, 2004.
- Ergonomics Guideline for Furniture used in Office Work Spaces Designed for Computer Use (2013). Business & Institutional Furniture Manufacturers Association.
- Hedge, A., personal communication, July 14, 2016.
- Hedge, A., (2015). Sit-Stand Working Programs. Retrieved October 3, 2016 from <http://ergo.human.cornell.edu/CUESitStandPrograms.html>
- Human Factors Engineering of Computer Workstations (2007). Human Factors and Ergonomics Society. ANSI/HFES 100-2007.
- Human Factors and Ergonomics Society. (2015, July 14). Like sitting, standing in the workplace may have long-term health consequences. *ScienceDaily*. Retrieved July 10, 2017 from www.sciencedaily.com/releases/2015/07/150714093611.htm
- Karakolis & Callaghan. (2014). The impact of sit–stand office workstations on worker discomfort and productivity: A review. *Applied Ergonomics*, 45(3), 799-806.
- Krijnen, R. M., De Boer, E. M., Adèr, H. J., Osinga, D. S., & Bruynzeel, D. P. (1997). Compression Stockings and Rubber Floor Mats: Do They Benefit Workers With Chronic Venous Insufficiency and a Standing Profession? *Journal of Occupational & Environmental Medicine*, 39(9), 889-894.

Konz, S., & Johnson, S. (2000). *Work Design: Industrial Ergonomics* (5th ed.). Scottsdale, Ariz.: Holcomb Hathaway.

Lin, M., Catalano, P., & Dennerlein, J. (2016). A Psychophysical Protocol to Develop Ergonomic Recommendations for Sitting and Standing Workstations. *Human Factors: The Journal of Human Factors and Ergonomics Society*, 58(4), 574-585.

Neuhaus, Healy, Dunstan, Owen, & Eakin. (2014). Workplace Sitting and Height-Adjustable Workstations: A Randomized Controlled Trial: A Randomized Controlled Trial. *American Journal of Preventive Medicine*, 46(1), 30-40.

Owen, N., Healy, G., Matthews, C., Dunstan, D., (2010) Too Much Sitting: The Population Health Science of Sedentary Behavior. *Exercise and Sport Sciences Reviews* 38: 105–113.

Pheasant, S. (1986). *Bodyspace: Anthropometry, ergonomics, and design*. London; Philadelphia: Taylor & Francis

Reinhardt, F., Wetzel, T., Vetten, S., Radespiel-Troger, M., Hilz, M. J., Heuss, D., & Neundorfer, B. (2000). Peripheral neuropathy in chronic venous insufficiency. *Muscle & Nerve*, 23, 883–887.

Sanders, M. S. (1993). *Human factors in engineering and design*. New York: McGraw-Hill.

Sandy, M., Mason, Thomas J., Bernard, Thomas E., Banks, Anthony, & Mylnarek, Steven. (2016). Longitudinal Study of Adjustable Workstations, ProQuest Dissertations and Theses.

Van Der Ploeg, Chey, Korda, Banks, & Bauman. (2012). Sitting time and all cause mortality risk in 222,497 Australian adults. *Journal of Science and Medicine in Sport*, 15, S28.

Vernikos, J., National Ergonomics Conference, personal communication, Nov 2015.

Wang, A., Arah, O., Kauhanen, J., & Krause, N. (2015). Work schedules and 11-year progression of carotid atherosclerosis in middle-aged Finnish men. *American Journal of Industrial Medicine*, 58(1), 1-13.

Wahlstrom, J. (2005). Ergonomics, musculoskeletal disorders and computer work. *Occupational Medicine-Oxford*, 55(3), 168-176.

Wilke, H. E., Neef, P., Caimi, M., Hoogland, T., & Claes, L. (1999). New In Vivo Measurements of Pressures in the Intervertebral Disc in Daily Life. *Spine*, 24(8), 755-762.

Wilmot, E., Edwardson, G., Achana, C., Davies, L., Gorely, F., Gray, A., Khunti, K., Yates, T., Biddle, T. (2012). Sedentary time in adults and the association with diabetes, cardiovascular disease and death: Systematic review and meta-analysis. *Diabetologia*, 55(11), 2895-2905.